


MASS OF CLUSTERS IN SIMULATIONS

A.V. MACCIÒ^{1,2}, G. MURANTE³, S.A. BONOMETTO^{1,2}

¹ *Dep. of Physics G. Occhialini, Milano Bicocca Univ., Piazza della Scienza 3, 20126 Milano (Italy)*

² *I.N.F.N. – Sezione di Milano*

³ *I.N.A.F., Osservatorio Astronomico di Torino - Torino (Italy)*



We show that dark matter haloes, in n-body simulations, have a *boundary layer* (BL) with precise features. In particular, it encloses all dynamically stable mass while, outside it, dynamical stability is lost soon. Particles can pass through such BL, which however acts as a confinement barrier for dynamical properties. BL is set by evaluating kinetic and potential energies ($T(r)$ and $W(r)$) and calculating $\mathcal{R} = -2T/W$. Then, on BL, $\mathcal{R}$ has a minimum which closely approaches a maximum of $w = -d \log W / d \log r$. Such Rw “requirement” is consistent with virial equilibrium, but implies further regularities. We test the presence of a BL around haloes in spatially flat CDM simulations, with or without cosmological constant. We find that the mass M_c , enclosed within the radius r_c , where the Rw requirement is fulfilled, closely approaches the mass M_{dyn} , evaluated from the velocities of all particles within r_c , according to the virial theorem. Using r_c we can then determine an individual density contrast Δ_c for each virialized halo, which can be compared with the “virial” density contrast $\Delta_v \simeq 178 \Omega_m^{0.45}$ (Ω_m : matter density parameter) obtained assuming a spherically symmetric and unperturbed fluctuation growth. The spread in Δ_c is wide, and cannot be neglected when global physical quantities related to the clusters are calculated, while the average Δ_c is $\sim 25\%$ smaller than the corresponding Δ_v ; moreover if M_{dyn} is defined from the radius linked to Δ_v , we have a much worse fit with particle mass then starting from Rw requirement.

1 Introduction

Finding the volume where cluster materials are in virial equilibrium might seem a tough and somewhat ambiguous task, as clusters are not surrounded by vacuum and shall be however confined by unbound background materials. Here we show that, in general, the transition from cluster and external materials is well defined and takes place just outside a *boundary layer* (BL), which, by itself, is in virial equilibrium with no pressure pushes, but above which virial equilibrium is lost soon. Particles may pass through such BL, in both directions, but its overall virial balance keeps constant in time. We find such BL assuming a spherical symmetry. At a distance where the cluster ends up into unbound materials, such requirement might seem excessive. Rather surprisingly, however, we shall see that this is not a prohibitive restriction.

Previous work on this point was based on the assumption of isolated fluctuation growth, starting from and preserving spherical symmetry (Gott & Rees 1975, Lahav et. al. 1991, Eke

et al. 1996, 1998, Brian & Norman 1998). Results obtained in this way were then applied to simulation analysis. On the contrary, in our work, the assumption of isolated growth is absent, while spherical symmetry is tested, *a posteriori*, on simulated clusters. In this way we find that actual clusters closely approach properties predicted under such assumption.

2 How to find BL

Let us define *integral* potential and kinetic energy and their *virial ratio*:

$$2T(< r) = \sum_i (r_i < r) m v_i^2, \quad W(< r) = - \sum_{i < j (r_{i,j} < r)} \frac{G m^2}{r_{ij}}, \quad \mathcal{R}(< r) = - \frac{2T(< r)}{W(< r)}. \quad (2.1)$$

Similar quantities can be defined for r -intervals. The r dependence of W can be outlined by performing the sums in eq. (2.1) starting from outside:

$$W(< r) = - \sum_i (r_i < r) \sum_j (r_j < r_i) \frac{G m^2}{r_{ij}} = - \sum_i (r_i < r) \mathcal{Z}(r_i). \quad (2.2)$$

Here the last term is the very definition of $\mathcal{Z}$. Quite in general, a volume integral of $\rho(r)$ increases with r . Henceforth, $\mathcal{Z}(r) = C(r/\bar{r})^{-w}$, where C is a normalization constant evaluated at an arbitrary position $\bar{r}$ and $w(< 1)$ shall however depend on r . Accordingly, it must be

$$r \frac{d\mathcal{Z}}{dr}(r) = -[w + r w' \ln(\frac{r}{\bar{r}})] \mathcal{Z}(r). \quad (2.4)$$

Let us now suppose that there exists an interval $\Delta r = r_+ - r_-$, (i) in virial equilibrium, (ii) subject to no pressure pushes inwards or outwards, (iii) inside which w is constant. Owing to the conditions (i) and (ii), in this layer

$$\sum_i (r_i \in \Delta r) m v_i^2 - r_i \frac{d\mathcal{Z}}{dr}(r_i) = 0. \quad (2.5)$$

Taking then into account the extra condition (iii), i.e. that w is constant, we can use eq. (2.4) with $w' = 0$ and the expression for the *virial ratio* in Δr to obtain that

$$\mathcal{R} = w, \quad \frac{d\mathcal{R}}{dr} = 0 \quad (2.6)$$

all along the interval Δr . *Viceversa*, if the eq.s (2.6) are both fulfilled in a layer of depth Δr , such layer is at rest and in virial equilibrium. We thus define *boundary layer (BL)* a region of depth Δr satisfying eq.s (2.6). It can also be shown that no further materials can be in virial equilibrium if we required that w is maximum in Δr and therefore $w' < 0$ for $r > r_+$.

3 Application to numerical simulations

We seek BL for clusters, in simulations. We use two main sets of simulations, performed with different codes. The former set comprises large volume simulations (box side = $360h^{-1}$ Mpc), run with a parallel AP3M code developed by Gardini et al. (1999). They describe a tilted CDM (TCDM) model ($n = 0.8$) and a Λ CDM model ($\Omega_m = 0.35$, $\Omega_\Lambda = 0.65$) and were also used in Macciò et al. (2001; simulations A and B respectively). This set of simulations is meant to provide a significant statistics, although their mass (180^3 particles) and force resolution (~ 40 kpc P.E.) is limited. The second set of simulations aims to provide high-resolution clusters, using

the PM ART code (Kravtsov, Klypin & Khokhlov, 1997) and/or the public parallel tree-code GADGET (Springel et. al 2001). Here we resolve clusters with more than 300.000 particles within a radius of $2 h^{-1}$ Mpc, with a mass resolution of $1.2 \times 10^{10} \Omega_m h^{-1} M_\odot$. With these codes we performed simulations C and D of Λ CDM ($\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$) and TCDM ($n = 0.8$) models, obtaining 6 high resolution clusters for each model.

To set the position of BL, in each dark matter halo, we first calculate the virial ratio $\mathcal{R}$ in successive layers; then, we select points where $\mathcal{R}$ has a minimum and (nearly) intersects w (so to fulfill eq.s 2.6). Such coincidence is shown in Fig. 1 for the best and worst cases we found in the simulations C and D. Even in worst case, the coincidence between a minimum of $\mathcal{R}$ and a maximum of w is neat. In simulations A and B, w is not so well traced, as it is obtained from differentiation, and is only used to select among minima of $\mathcal{R}$, when we have several of them. The procedure however works in 97 % of cases, allowing to find a precise BL.

Once the sphere confining cluster materials is set, we can evaluate the density contrast Δ_c and the mass M_c therein. In Figs 2 and 3, points give Δ_c and M_c for all clusters in simulations A and B, respectively. They show that the spread of Δ_c values is fairly wide. By subdividing the M_c abscissa in intervals of constant logarithmic width, we evaluate the average density contrast in each of them, to seek systematic trends with mass.

Owing to the spread of Δ_c values, their averages are still subject to a significant uncertainty, shown, at the $1-\sigma$ level, in the plots. There seems to be no evidence of any peculiar trend of density contrasts with mass apart, perhaps, a modest indication of an increasing density contrast, at very high scales, in Λ CDM. It is therefore licit to consider the overall average among Δ_c 's. Such average is indicated by the continuous horizontal line and compared with the "virial" density contrast Δ_v , as given by eq. (1.1). In both cases, Δ_v (dotted line) is well inside the range of the density contrasts we found; the average Δ_c , however, in both cases, is smaller than Δ_v by $\sim 25\%$.

A critical result of our analysis is however shown in Figs 4 and 5 (for simulations A and B, respectively). In simulations, cluster masses can be comfortably evaluated by summing up particle masses and we compared this mass estimate with $M_{dyn} = \frac{\langle v^2 \rangle_{\{v,c\}} r_{\{v,c\}}}{G}$, evaluated averaging over the velocities of the particles within $r_{v,c}$ (r_v is the radius whereinside the density contrast is Δ_v). This comparison yields a reasonable coincidence between M_v and M_{dyn} as is shown in Figs 4 and 5 by the dashed histograms (which confirm the slight excess of M_{dyn} vs. M_v already noticed by previous authors). In the same plots we also report the much better fit between M_{dyn} and M_c , obtained on the basis of the setting of the BL (continuous line). The average values of M_{dyn}/M_c are $\sim 0.97 \pm 0.03$ for both models.

4 Conclusions

Our results are based on the mathematical definition of a boundary layer (BL), bordering a region satisfying the virial theorem, although its density contrast has different values in different haloes. The BL is found in numerical simulation, around each cluster, and it is shown to separate bound cluster materials from the surrounding medium. Boundary layers can be found in simulations with sufficient dynamical range, and have been confirmed by inspections of higher resolution simulations. The spread of the density contrasts of individual clusters has been found to be fairly wide and, in average, Δ_c are smaller by $\sim 25\%$ when compared to the so-called "virial" density contrast ($\Delta_v \simeq 178 \Omega_m^{0.45}$). No evidence of a systematic trend of Δ_c with M_c was found. The fit between dynamical masses and particle masses, obtained with our technique, is however far better than the one obtained starting from a fixed virial density contrast.




Fig. 1




Fig.2




Fig.3




Fig 4




Fig.5

Acknowledgments

We thank INAF for allowing us the CPU time to perform the ART simulation C and D at the CINECA consortium (grant cnami44a on the SGI Origin 3800 machine). GADGET simulations of high-resolution clusters have been run on the 16 Linux PC Beowulf cluster at the Osservatorio Astronomico di Torino.

References

1. Brian, G. & Norman, M., 1998, ApJ, 495, 80
2. Eke, V.R., Cole, S., & Frenk, C.S., 1996, MNRAS, 282, 263
3. Eke, V., Navarro, J. & Frenk, C., 1998, ApJ, 503, 569E
4. Gardini, A., Murante, G. & Bonometto, S. A., 1999, ApJ, 524, 510
5. Gott, R. & Rees, M. 1975, A&A, 45, 365G
6. Kravtsov A., Klypin A. & Khokhlov A., 1997 ApJ, 111, 73 K
7. Lahav, O., Lilje, P. R., Primack, J. R. & Rees, M., 1991, MNRAS, 282, 263E
8. Macciò A. V., Gardini A., Ghigna S. & Bonometto S. A., 2002, ApJ, 564, 1M
9. Peebles P.J.E., 1980, The Large Scale Structure of the Universe, Princeton University Press, Princeton
10. Springel V., Yoshida N. & White S.D.M., 2001 NewA, 6, 79 S